

This datasheet offers general information. Please contact your local ECO representative for country-specific details.



ECOTRAZ PLUS POUR-ON

Presentation

Ready-to-use pour-on for cattle, sheep and goats that contains 1.5% m/v amitraz, 0.5% m/v deltamethrin & 3.0% m/v piperonyl butoxide.

Indications

Cattle

For the control of ticks, lice (biting and sucking), mites, stable flies, horn flies, cattle louse flies and nuisance flies, e.g. houseflies. Protects against black flies (*Simulium* spp).

Sheep and goats

For the control of ticks including paralysis ticks, bont-legged (*Hyalomma* spp) and red-legged ticks (*Rhipicephalus evertsi*).

Dosage and administration

Cattle

The dose is 1ml/10kg bodyweight. Apply in two lines from the base of the skull down both sides of the spine to the tail root. Application at the base of the skull is important in the case of heavy ear tick infestation. Treat at 7 day intervals according to tick challenge. The longer the product is applied at regular intervals, the more the tick population is reduced. It may then become possible to apply at intervals greater than 7 days.

Sheep and goats

Dosage for optimal residual activity is 1ml/5kg bodyweight. Apply to the axillae and groin region, whilst the animal is turned over in a sitting position. Treat at intervals of 2-4 weeks, according to tick infestation.

Contra-indications, warnings etc.

Withdrawal period

- Meat – 7 days
- Milk - nil
- Do not use on calves or lambs under 3 months of age.
- Treat strictly according to body weight.

Minor signs of discomfort have been noted in some cattle during the 48 hours after treatment; this is of no long-term detriment to the animal.

This datasheet offers general information. Please contact your local ECO representative for country-specific details.

Operator warnings

- Wear overalls, rubber gloves and face shield when opening containers or diluting, mixing or transferring the product from one container to another or when washing out the container.
- Wear arm-length rubber gloves, rubber boots and bib apron when handling diluted fluid or freshly treated animals.
- When using do not eat, drink or smoke.
- Do not breathe spray mist.
- Avoid all contact with mouth, skin and eyes.
- Wash concentrate or splashes of diluted spray from skin or eyes immediately.
- Wash all protective clothing thoroughly after use especially the inside of gloves.
- Wash hands and exposed skin before eating, drinking or smoking and after work.
- If you feel unwell seek medical advice and show this document.
- Containers should be safely disposed of after use and not re-used.
- Do not store in residential properties.

Guide to Doctor

- Amitraz is **NOT** an organophosphate compound. If poisoning is suspected treatment should be symptomatic and supportive, paying particular attention to monitoring of respiratory and cardiac function.
- Do **NOT** use Atropine.
- Do **NOT** induce vomiting.

Protection of wildlife and the environment

Harmful to fish. Do not contaminate ponds, waterways or ditches with chemical or used container.

Hazard warnings

May cause sensitisation by skin contact. Harmful if swallowed.

Pharmaceutical precautions

- Keep away from food, drink and animal feedingstuffs.
- Keep out of the reach and sight of children and unauthorised persons.
- For animal treatment only.
- For external use only.
- Keep away from domestic animals.
- Store below 25°C in a dry place and protect from freezing.
- Shake well before use. Store in a cool, dry place.
- Close container after use.

General precautions

- Do not use on horses, dogs or cats.

Package quantities

1 litre & 5 litre containers

This datasheet offers general information. Please contact your local ECO representative for country-specific details.

Further information

For further information, contact:

ECO Animal Health
PO Box 47542
London N14 6WS
United Kingdom

Tel: +44 (0)208 447 8899
Fax: +44 (0)208 447 9292
Website: www.ecoanimalhealth.com