

This datasheet offers general information. Please contact your local ECO representative for country-specific details.



ECOTEL PLUS DRENCH

Presentation

Ecotel Plus Drench is an oral anthelmintic containing levamisole hydrochloride 37.5mg/ml and praziquantel 18.8mg/ml.

Indications

Ecotel Plus Drench is indicated as a roundworm, lungworm and tapeworm remedy for cattle, sheep, goats and ostriches.

Dosage and administration

Ecotel Plus Drench is recommended at the following dose rates:

Cattle: 1 ml/5 kg bodyweight
Sheep and Goats: 1 ml/5 kg bodyweight
Ostriches: 1 ml/1 kg bodyweight

Contra-indications, warnings etc

Withdrawal periods

- Do not slaughter animals for human consumption within 7 days of the last treatment.

Operator warnings

- Protective clothing must be worn when handling the product.
- When using do not eat, drink or smoke.
- Avoid all contact with mouth, skin and eyes.
- Wash splashes from skin or eyes immediately.
- Wash all protective clothing thoroughly after use especially the inside of gloves.
- If you feel unwell seek medical advice and show this document.

Protection of wildlife and the environment

Do not contaminate ponds, waterways or ditches with chemical or used container.

Hazard warnings

May cause sensitisation by skin contact. Harmful if swallowed.

Pharmaceutical precautions

- Store in original container. Ensure container is sealed securely after each use.
- Store below 25°C in a cool, dry place and protect from sunlight.

This datasheet offers general information. Please contact your local ECO representative for country-specific details.

- Keep away from food, drink and animal feedingstuffs.
- Keep out of the reach and sight of children.
- For animal treatment only.

Container disposal

Do not dispose of undiluted chemicals on site. Rinsed containers should be crushed or have holes put in and then disposed of in accordance with any guidance from an appropriate waste regulation authority. Unused product should be returned to a pharmacist for disposal.

General precautions

- No worm remedy should be dosed to sheep and goats that have not been immunised against pulpy kidney (enterotoxaemia).
- Levamisole is not effective against inhibited *Ostertagia* larvae in cattle.
- When treated with levamisole, some animals may develop an allergic-type reaction of a transient nature, which normally disappears within 24 hours. An antihistamine may be administered. Under no circumstances must cortisone preparations be used

Package quantities

Ecotel Plus Drench is supplied in 200ml, 1L, 5L and 10L containers.

Further information

For further information contact:

ECO Animal Health
PO Box 47542
London N14 6WS
United Kingdom

Tel: +44 (0)208 447 8899
Fax: +44 (0)208 447 9292
Website: www.ecoanimalhealth.com