

This datasheet offers general information. Please contact your local ECO representative for country-specific details.



LEVECO 2.5% DRENCH

Presentation

Leveco 2.5% is a colourless to pale red solution containing 2.5% w/v levamisole hydrochloride.

Indications

The product is indicated as a roundworm and lungworm remedy for cattle, sheep, goats and ostriches.

For the treatment of mature and developing immature stages of levamisole-susceptible nematode worm species.

- In the abomasum: *Haemonchus* (Wireworm, barberpole worm), *Ostertagia* (brown stomach worm), *Trichostrongylus spp.* (bankrupt worm, small stomach worm).
- In the intestines: *Trichostrongylus*, *Cooperia spp.* (small intestinal worm), *Nematodirus* (thread-necked intestinal worm), *Oesophagostomum* (nodular worm), *Chabertia* (large mouth bowel worm) and *Bunostomum spp.* (Hookworm).
- In the lungs: *Dictyocaulus spp.* (Lungworm).

Also effective for the treatment of *Libostrongylus douglassi* (Ostrich wireworm).

This anthelmintic may also be effective against other species. Consult your veterinarian.

Dosage and administration

Given as an oral drench. Shake well before use and use only as directed. Does not stain wool or mohair.

For cattle, sheep and goats, dose according to bodyweight at the rate of 7.5mg levamisole hydrochloride per kg bodyweight.

For Ostriches, dose according to bodyweight at a rate of 30mg levamisole hydrochloride per kg bodyweight.

Cattle: 15 ml/50 kg bodyweight
Sheep and Goats: 3 ml/10 kg bodyweight

This datasheet offers general information. Please contact your local ECO representative for country-specific details.

Ostriches: 12 ml/10 kg bodyweight

Example for cattle, sheep and goats:

Bodyweight	Dose	Number of Doses in Pack					
		200ml	250ml	500ml	1 L	5 L	10 L
10 kg	3 ml	66	83	166	333	1,666	3,333
20 kg	6 ml	33	41	83	166	833	1,666
30 kg	9 ml	22	27	55	111	555	1,111
40 kg	12 ml	16	20	41	83	416	833
50 kg	15 ml	13	16	33	66	333	666
100 kg	30 ml	6	8	16	33	166	333

Contra-indications, warnings etc

Withdrawal periods

- Cattle must not be treated within a period of 14 days before or after treatment with organophosphorus compounds.
- Do not slaughter animals for human consumption within 7 days of the last treatment.

Operator warnings

- Protective clothing must be worn when handling the product.
- When using do not eat, drink or smoke.
- Avoid all contact with mouth, skin and eyes.
- Wash splashes from skin or eyes immediately.
- Wash all protective clothing thoroughly after use especially the inside of gloves.
- Remove contaminated clothing and wash all exposed areas after use.
- If you feel unwell seek medical advice and show this document.

Protection of wildlife and the environment

Do not contaminate ponds, waterways or ditches with chemical or used container.

Hazard warnings

May cause sensitisation by skin contact. Harmful if swallowed.

This datasheet offers general information. Please contact your local ECO representative for country-specific details.

Pharmaceutical precautions

- Close container after use. Store in original container.
- Store below 25°C in a cool, dry place and protect from sunlight.
- Keep away from food, drink and animal feedingstuffs.
- Keep out of the reach and sight of children.
- For animal treatment only.
- For external use only.
- Keep away from domestic animals.

General precautions

- No worm remedy should be dosed to sheep and goats that have not been immunised against pulpy kidney (enterotoxaemia).

Package quantities

Leveco 2.5% is supplied in 200ml, 250ml, 500ml, 1L, 5L and 10L multi-dose containers.

Further information

For further information contact:

ECO Animal Health
PO Box 47542
London N14 6WS
United Kingdom

Tel: +44 (0)208 447 8899
Fax: +44 (0)208 447 9292
Website: www.ecoanimalhealth.com